

## Racial classification of the World:

Earnest Albert Hooton's classification. The most recent attempt is a debilitated phylogenetic (मूलवृक्ष) classification is that of Hooton. His efforts of classification are based on purely anatomical (आन्तरिक) characteristics:

### A. White, or Caucasoid Primary Race

#### 1. Mediterranean

a) Upper palaeolithic survivals, chiefly in British Isles.

b) Iranian plateau

c) Classic Mediterranean

#### 2) Ainu

3) Keltic (light eyed, dark or red haired, mainly in British Isles).

4) Nordic

5) Alpine

6) East Baltic

7) Armenoid

8) Dinaric

9) Nordic - Alpine

10) Nordic - Mediterranean

composite, predominantly white

#### 11) Australian

a) Murray, mostly nearly white, specially in south east.

b) Carpenterian, Melanesian increased in the North

c) Tasmanoid, refuge areas in Queensland

12) Indo - Dravidian (classic Mediterranean Australoid, Negrito)



- a) classic Indo-Dravidian, mostly in northern India.
  - b) Armenoid - Iranian plateau, (western and southern India)
  - c) Indo - Nordic North - Western Himalayas
  - d) Australoid or Veddoid - central and southern India.
  - e) Negritoid - spots in southern India
- 12) Polynesian.

### B) Negroid Primary Race

- 14) African Negro or forest Negro
- 15) Nilotic Negro
- 16) Negrito

- a) Infantile type, in all Negrito populations.
- b) Adultiform among all negritos, except in the Andamans and Philippines.

Composite, predominately Negroid

- 17) Tasmanian: Negrito plus Australians
- 18) Melanesian: Papuan, Negrito Australoid
  - a) Papuan
  - b) Melanesian
- 19) Bushmen - Hottentots
  - a) Bushmen
  - b) Hottentot, Bushman plus Negro

### C) Mongoloid Primary Race

- 20) classic Mongoloid
- 21) Asiatic Mongoloid or Eskimoid
- 22) Indonesian
  - a) Malay-Mongoloid (Indonesia and further north including most of the Japanese).



b) Lada region (pre-Mongoloid groups in south China, further in India)

23) American Indians

- a) Brachycephals,
- b) Dolichocephals.