

TYPES OF NEPHRIDIA :-

Nephridia are categorized into several types based on certain features.

- (A) Based on the tubes are closed or open.
- (B) Based on size and number of Nephridia.
- (C) Based on site of opening of Nephridiopore.

(A) Based on the tubes are closed or open:-

Based on the tube which are closed or open, Nephridia are divided into two types : - (i) Protonephridia and (ii) Metanephridia.

(i) Protonephridia : - The protonephridia are very primitive type of Nephridia. This nephridia terminates the coelom as blind tube. i.e. closed. The typical protonephridia terminates ~~with~~ ^{with} tube ¹ and specially excretory ~~some~~ ¹ cells ~~of~~ ¹ Flame Cells or Solenocytes. Protonephridia are found in ~~Phy~~ Phyllocladus, Vannadis Nephtys, Glycera etc.

protonephridia are considered as the most primitive form of nephridia basically these are ^{the} excretory organs of larvae and other adult polychaetes. They have a tubular arrangement that open outside the body by Nepridiopore. They have tubes with blind ends where cilia mediated filtration occurs. They have terminal or captrines which can be more or multi ciliated that support in filtration process. Protonephridia

are divided into two types based on their termination. These are flame cells or tube cell and Solenocytes.


(i) Flame cells:- The protonephridia which whose blind ends terminate in ~~fan~~ flame cells. These cells are bulb-like in appearance hence called flame bulb. These cells bears a cluster of cilia within a cup like projection whose beating is like a candle flame. A flame cell is a specialized cell. It consist of the extracellular processes, ciliary flame and intercellular duct with certain secretory materials. The intercellular duct joins with the excretory duct.

(ii) SOLENOCYTES :- The second type of protonephridia are the Solenocytes. Each nephridia bears about 500 solenocytes. These are flagellated, tiny rounded nucleated cell body and a long & hollow stalk or

tubules. Solenocytes can be distinguished from flame cell as the former is usually flagellated whereas the latter is ciliated. The solenocytes bear a flagella whereas the flame cell bears a part of cilia. The solenocytes associated with excretion, osmoregulation and ion regulation in many annelids. These are found in Phyllodoce and other members of Annelids.

Metanephridia — These are far advanced nephridia found in vast majority of Annelids. It is a nephridia that originated in a ciliated coelomic funnel. In metanephridia the inner of the tube is open and remain in the form of a nephrostom. A typical metanephridia consist of a nephrostom a short co-

^ Protonephridia



Flatworm flame cell

A **protonephridium** (*proto* = "first") is a network of dead-end **tubules** lacking internal openings, found in the phyla **Platyhelminthes**, **Nemertea**, **Rotifera** and **Chordata** (lancelets). The ends are called *flame cells* (if ciliated) or *solenocytes* (if flagellated); they function in **osmoregulation** (ionoregulation). The terminal cells are located at the blind end of the




Fig. 8.36. Protonephridium and coelomoduct in *Phyllocoete*. **A**—Protonephridium with coelomoduct, **B**—Solenocytes of one protonephridium, **C**—Group of solenocytes.




Fig.- Showing Protonephridium of *Phyllodoce paretti*
 (From M.Raj & Hazarika (2020); Barnes (1987))

cells




Fig : A Flame Cell
(From <https://seunojo.blogspot.com>)

Cannal and a nephri duct and opens through a nephridiopore to the exterior.

Metanephridia found in oligochaetes (*Lumbricus*), polychaetes (*Nereis*), Leeches etc.

(B) Based on size and ^{number} of Nephridia :— The second category of nephridia are based on the ~~number~~

number of nephridia per segment are categorized into (1) Micro nephridia and ~~Mega nephridia~~ or Mesonephridia and (2) Mega nephridia or Hollow nephridia.

(1) Micro nephridia — Smaller in size, but numerous per segment. The network of fine tubes lie lying on the body wall and septa in each segment. The nephridia of *Pheretima postuma* are micro nephridia type.

The

(2) Meganephridia - True meganephridia are larger in size but generally one pair per segment. ~~at least~~

Usually they extend over two segments and their nephrostomes open into the next segment.

The Nepridinia of Hirudinaria are the examples of mega nephridia.

(c) Based on Site of opening :-

The third category of Nephridia are based on site of opening. Based on the site of opening of the Nepridiosome the nephridia are classified in two types:

(i) Exo-nephridia or Ectonephridia

(ii) Entero nephridia

(i) Exonephridia -

The Exonephridia directly ~~open~~ opens to the exterior. Nephridiopore are present.

In case of Exonephridia. The integumentary micronephridia in ~~Fec~~ pherotima postuma, the meganephridia of ~~adonis~~, Hirudin Lumbriens etc. are the example of Exonephridia.

(ii)

Entero nephridia - Entero nephridia opens to the excretory canal or alimentary canals. Nephridiopore absent. The Septal and pherangial nephridia of pherotima postuma are the example of Entero nephridia.

chloragogen Cells:- These cells are star-shaped and are also called as γ -cells. These are derived from inner coelomic epithelium with excretory function. These cells are found in earth worm.

They are yellowish in color due to yellow granules chloragosomes.

The cell stores glycogen and neutralizes toxins ⁱⁿ and are present in the coelomic fluid of some Annelids.

These cells function as liver in annelids and store and transport

they store and transport nitrogenous waste like urea.

nephridia




Fig : Metanephridium of the night-crawler earthworm *Lumbricus*